

10



Based on Latest CBSE Syllabus

LIVING SCIENCE LAB MANUAL

With Practical Skills


Ratna Sagar

Based on the latest CBSE syllabus and guidelines

LIVING SCIENCE LAB MANUAL

With Practical Skills

10



HITI MAHINDROO



Ratna Sagar

ABOUT THE AUTHOR

Hiti Mahindroo completed BSc (Physics Hons) and MSc (Physics) from the Physics Honours School, Panjab University, Chandigarh in 1972. She has taught Physics for many years, both at college and high school levels, and Science at middle school level.

© Ratna Sagar P. Ltd., 2012
All rights reserved
First published 2012
Seventh reprint 2017 (March)

ISBN 978-93-5036-152-8



Ratna Sagar P. Ltd.

an ISO 9001:2015 and 14001:2015 company

VIRAT BHAVAN, COMMERCIAL COMPLEX, MUKHERJEE NAGAR, DELHI 110009

PHONE: (011) 47038000 ♦ FAX: (011) 47038099

rsagar@ratnasagar.com ♦ ratnasagar.com ♦ ratnasagar.co.in

SHOWROOM: 4808/24 BHARAT RAM ROAD, DARYA GANJ, NEW DELHI 110002

PHONE: (011) 43028000 ♦ FAX: (011) 45166099

BRANCHES: CHENNAI ♦ LUCKNOW

OFFICES: AGRA ♦ AHMADABAD ♦ BENGALURU ♦ BHOPAL ♦ COIMBATORE ♦ DEHRADUN
GUWAHATI ♦ HYDERABAD ♦ JAIPUR ♦ JALANDHAR ♦ KANPUR ♦ KOCHI ♦ KOLKATA
MADURAI ♦ MUMBAI ♦ PATNA ♦ RANCHI ♦ VARANASI

Contents

In the Lab	5
1. pH of Solutions	15
Experiment 1: To find the pH of the following samples by using pH paper/universal indicator. 1. Dilute hydrochloric acid (HCl) 2. Dilute sodium hydroxide (NaOH) solution 3. Dilute ethanoic acid (CH ₃ COOH) solution 4. Lemon juice 5. Water (H ₂ O) 6. Dilute sodium bicarbonate (NaHCO ₃) solution.	
2. Properties of Acids and Bases	24
Experiment 2A: To study the properties of an acid (dilute HCl) by its reactions with 1. Litmus solutions (blue/red) 2. Zinc metal 3. Solid sodium carbonate Experiment 2B: To study the properties of a base (dilute solution of NaOH) by its reaction with 1. Litmus solutions (blue/red) 2. Zinc metal 3. Solid sodium carbonate	
3. Combination, Decomposition, Displacement and Double Decomposition Reactions	35
Experiment 3A: To perform and observe the action of water on quicklime (calcium oxide); and classify it as a combination reaction. Experiment 3B: To perform and observe the reaction caused by action of heat on iron(II) sulphate (ferrous sulphate) crystals, and to classify it as a decomposition reaction. Experiment 3C: To perform and observe the reaction between iron nails kept in copper sulphate solution and to classify it as a displacement reaction. Experiment 3D: To perform and observe the reaction between sodium sulphate and barium chloride solutions, and classify it as double displacement reaction.	
4. Reactivity of Metals	50
Experiment 4: a. To observe the action of zinc (Zn), iron (Fe), copper (Cu) and aluminium (Al) metals on the following salt solutions: i. Zinc sulphate solution [ZnSO ₄ (aq)] ii. Ferrous sulphate solution [FeSO ₄ (aq)] iii. Copper sulphate solution [CuSO ₄ (aq)] iv. Aluminium sulphate solution [Al ₂ (SO ₄) ₃ (aq)] b. To arrange zinc (Zn), iron (Fe), copper (Cu) and aluminium (Al) metals in the decreasing order of their reactivity, based on the above result.	
5. Electric Current, Potential Difference and Resistance	63
Experiment 5: To study the dependence of potential difference (<i>V</i>) across a resistor on the current (<i>I</i>) passing through it and to determine its resistance. Also plot a graph between <i>V</i> and <i>I</i> .	
6. Resistances in Series	74
Experiment 6: To determine the equivalent resistance of two resistors when connected in series.	
7. Resistances in Parallel	81
Experiment 7: To determine the equivalent resistance of two resistors when connected in parallel.	
8. Stomata on Leaf Surface	89
Experiment 8: To prepare a temporary mount of a leaf peel to show its stomata.	

9. Photosynthesis	96
Experiment 9: To show experimentally that light is necessary for photosynthesis.	
10. Respiration	103
Experiment 10: To show experimentally that carbon dioxide is given out during respiration.	
11. Acetic Acid (Ethanoic Acid)	111
Experiment 11: To study the following properties of acetic acid (ethanoic acid): 1. Odour 2. Solubility in water 3. Effect on litmus 4. Reaction with sodium bicarbonate.	
12. Saponification: The Process of Making Soap	118
Experiment 12: To study the saponification reaction for preparation of soap.	
13. Cleaning Capacity of Soap with Hard and Soft Water	125
Experiment 13: To study the comparative cleaning capacity of a sample of soap in soft and hard water.	
14. Reflection by Concave Mirror	131
Experiment 14: To determine the focal length of a concave mirror by obtaining the image of a distant object.	
15. Refraction through a Convex Lens	137
Experiment 15: To determine the focal length of a convex lens by obtaining an image of a distant object.	
16. Refraction through a Glass Slab	143
Experiment 16: To trace the path of a ray of light through a rectangular glass slab for different angles of incidence. Measure the angle of incidence, angle of refraction and angle of emergence and interpret the result.	
17. Binary Fission and Budding	151
Experiment 17: To study A. binary fission in <i>Amoeba</i> and B. budding in yeast with the help of prepared slides.	
18. Images Formed by a Convex Lens	158
Experiment 18: To find the image distance for varying object distances in case of a convex lens and draw corresponding ray diagrams to show the nature of image formed.	
19. Prisms	167
Experiment 19: To study the path of a ray of light and find its deviation, when it passes through the prism.	
20. Seed Structure and Germination	175
Experiment 20: To identify the different parts of an embryo of a dicot seed (pea, gram or red kidney beans).	
21. Homology and Analogy in Plants and Animals	181
Experiment 21A: To identify and study homologous organs in animals and plants.	
Experiment 21B: To identify and study analogous organs in animals and plants.	
Test Papers	190

In the Lab

GENERAL LABORATORY FACILITIES

A science laboratory is equipped with working tables, items of common utility and space for storing equipment, chemicals and glassware.

1. The working tables in a science laboratory are usually fitted with the following:
 - Sinks with water taps for washing purposes and liquid waste disposal. Students are expected to use taps only when required and will not waste water.
 - Heating facilities provided in the form of gas taps fitted with a burner. Leakage, if any, should be immediately brought to the notice of the teacher. Also, spirit lamps can be provided.
 - Shelves for keeping bottles of chemicals and reagents of frequent use. These reagent bottles should be arranged in a definite order.
2. Equipment and glassware of common use are stored separately in a cupboard. They are generally issued to the students at the time of performing experiments.
3. Usually, instruments which are not used quite frequently, like balances and microscopes are kept permanently in a place. The type of balance used depends on how much accuracy is required of the weighing. At secondary stage, a physical balance is a good choice.

The microscopes must be placed near the window to ensure the availability of sufficient sunlight needed.

4. Electrical connections must be provided with all working tables for performing electricity and magnetism experiments.
5. A demonstration table is necessary for teacher's demonstration.
6. For the purpose of disposal of solid waste, a dustbin may be provided either near each working table or at a common place in the laboratory.

General Laboratory Rules

The following rules should be observed by all users in a science laboratory:

1. Students must prepare in advance for the experiment they would perform in the laboratory.




Fig. 1 A physical balance

2. Students must carry their practical notebook, laboratory manual, geometry box, pencil, sharpener, and eraser to the practical class regularly. They should also wear laboratory coats while working in the laboratory.
3. All equipment such as microscope, physical balance, measuring items and glassware must be handled with care after understanding their functions.
4. The labels on the reagent bottles should be read carefully to identify the reagents needed. To prevent contamination of reagents, do not dip glass rods, filter papers, droppers, etc. into the reagent bottles. Partially used materials should not be poured back into the reagent bottles. The cap/stopper should be put back on the bottle immediately after using the chemical reagent.
5. Windows and doors should be kept open.
6. Students should keep working tables clean and replace all laboratory equipment to their assigned place in the laboratory once the practical work is over.
7. Students should always wash their hands after the laboratory class.
8. Students should be acquainted with the handling of first-aid kit and fire extinguisher.
9. In case of any accident or injury or breakage of apparatus, report to the teacher immediately.

SAFETY REQUIREMENTS IN A SCIENCE LABORATORY

The following safety requirements are necessary in a science laboratory.

1. The laboratory should be well-ventilated. Provision for fuming hood or exhaust may be made in the laboratory to expel hazardous gases.
2. A first-aid box must be readily available, and the science teacher/lab assistant should be well versed with its use.
3. It is essential that every laboratory be equipped with a few fire extinguishers fixed at convenient places. Sand or dry mud should be available, (usually this is provided in buckets).
4. An *eye wash fountain* should be available in the laboratory. It is used in the case of irritation of eyes caused by chemicals and fumes.

First-Aid Treatment

A first-aid kit is an essential part of any science laboratory. In case of an accident, first-aid treatment must be provided immediately. The victim should be taken to the doctor, if required. Some common injuries caused due to accidents and their first-aid treatments are as follows:

Burns: In case of burns following treatment can be provided.

- For burns caused by steam or hot water, wash the affected part with cold running water. Do not use ice. Dry it gently with clean cloth and then apply an antiseptic ointment.
- For an acid burn, wash the wounded part with water and apply a dilute solution of sodium hydrogen carbonate. Dry gently with clean cloth and apply an antiseptic ointment.
- For an alkali burn, wash with a good amount of water to remove all the alkalis and then wash with dilute acetic acid. Dry the affected part with clean cloth and apply an antiseptic ointment.

Glass Cuts and Wounds: Carefully remove all the visible pieces of glass from the wound. Wash with cold water to remove smaller pieces of glass sticking to the wound, if any. Protect the wound from dirt and dust. Control the bleeding by pressing clean cotton or cloth on the wound. Apply an antiseptic solution/antiseptic cream. If required, take the victim to the doctor.

Eye Injuries: In case of injury to eyes, wash the eyes with cold water. Do not rub the eyes. Consult the doctor immediately.

Fires: Do not run around if clothes catch fire. Instead, lie down on the floor and roll. If a container with inflammable liquid catches fire while heating, turn off the gas burner immediately. Cover the mouth of the container with damp cloth and take it away from other reagents and chemicals. Fire extinguishers should be used if the fire goes out of control. In case of the short-circuit in electrical circuits, switch off the main switch of the electric supply and throw dry soil in the affected area. *Do not use water in case of electrical fire.*

Inhalation of Gases: If gases such as sulphur dioxide, chlorine or bromine are inhaled by anyone, take him/her into the open air immediately, and let the victim breathe deeply.

IN THE PHYSICS LABORATORY

General Precautions while doing Physics Experiments

Some special precautions should be followed while doing experiments in Physics.

1. *Percentage error* between 5% to 10% for recording observations and calculation of result is allowed. Percentage error is calculated by the formula:

$$\text{Percentage error} = \frac{\text{Result} - \text{Standard Result}}{\text{Standard Result}} \times 100$$

2. *Error of parallax* should be avoided by placing your eye at right angle to the plane of the scale, and vertically above the point to be read.
3. *Error due to spilling* arises when a solid or liquid is added to another liquid in a hurry. To avoid this error, a solid or liquid is added to another liquid gently, so that spilling does not occur.
4. *Error due to friction* may arise if movement of an object over another surface is involved. Hence, the error due to friction should be minimised. For example, in a spring balance, the strip carrying the pointer should be lubricated, and tapped gently before use.
5. *Error due to suspension* can arise if the support from which an object is suspended moves during the oscillation of the suspended object, i.e. in the case of a simple pendulum. To avoid this error, the support should be rigid, so that time period can be measured accurately.
6. *Error due to loss of heat* may arise in heat experiments, if a hot object is not transferred quickly to a calorimeter. To avoid this error, the calorimeter should be well insulated, and its contents should be stirred well with the stirrer, and the hot object should be transferred quickly to it.
7. *Accuracy* of a measuring instrument is taken as half of its least count. *Least count* is the smallest distance that measuring instruments like scales, Vernier callipers, screw gauge, spring balance, thermometer, etc. are able to measure.
8. *Zero error* arises in a measuring instrument when its pointer, at its resting stage, does not coincide with the zero mark on the scale. To obtain the accurate value, the zero error has to be added or subtracted from the reading taken after measurement.

PLOTTING A GRAPH

A graph shows the interdependence of two physical quantities. The quantity which is made to vary at will is called an *independent variable*, whereas the quantity that varies as a result is called a *dependent variable*. A graph is either a straight line or a curve.

Uses of a graph

1. A straight line graph shows that both the variables are directly proportional to each other.
2. Elaborate mathematical calculations are not required to derive the mean value of the ratio of the two variables.
3. The probable value of the variables can be calculated from the graph, even beyond the range of observations taken.

Steps to plot a graph

1. *Choosing axes:* If all data to be plotted is positive, draw two straight lines starting from the bottom left-hand corner of the graph paper, one horizontal line (X-axis) and the other a vertical line (Y-axis). The point of intersection of the axes is called origin, and is labelled O. If data is partly negative, then the origin O is shifted towards the middle of the graph paper.
2. *Which variable on which axis:* The independent variable is plotted always on X-axis, whereas the dependent variable is plotted on Y-axis.
3. *Choosing a suitable scale:* Choose any convenient scale to represent the variables, such that whole range of observations taken are well spread out on the whole graph paper as shown in Graph 2a). It is not necessary to take the origin from (0, 0), unless stated by the examiner.

It is not necessary that the scales on both X and Y axes should be same. Represent the axes by drawing thick lines at right angles to each other, beginning from the bottom left-hand corner of the graph paper. Mark each with an arrowhead.

4. *Denoting the scale chosen:* On top of graph paper, write down the scale chosen for X-axis and Y-axis. Mark the scale ratio (mentioning the proper units) taken along the X-axis and Y-axis.
5. *Marking the points:* Use a sharp pencil to mark the points of intersection. Encircle each point with a small circle (O).
6. *Plotting of graph:* It is not necessary that the graph line should pass through all the marked points. The line should pass through a maximum of marked points, or it should be plotted in such a way that it passes closest to maximum points on either side of it. To plot a curve, draw a smooth freehand curve which passes through maximum number of points.




Fig. 2a Correct selection of scale for plotting the graph




Fig. 2b Incorrect selection of scale for plotting the graph




Fig. 3 Accessories used in Biology laboratory

IN THE BIOLOGY LABORATORY

A dissection box

A dissection box should always be carried by students to the Biology laboratory. It should contain a pair of small forceps, a pair of large forceps, a pair of small scissors, a pair of large scissors, a fine razor for section cutting, a scalpel, a fine needle, two fine mounted needles, a fine brush and a magnifying lens.

To Make a Temporary Wet Slide (Mount)

Steps to make a temporary wet slide are given below.

- a. *Staining the specimen:* The specimen often requires staining, for better observation. Different stains such as safranin, iodine solution, methylene blue, cotton blue, etc. can be used for this purpose. Take a little stain in a watch glass. Transfer the specimen to be stained to it and allow to remain in the stain for a while. Remove the specimen from the stain and put it in another watch glass containing clean water in order to wash off extra stain.
- b. *How to proceed:* Hold a clean glass slide by its edges. Find the mid-point of the slide by drawing its diagonals on a rough sketch of the slide. Point of intersection of diagonals gives the mid-point of the slide. Put a drop of a clean water in the middle of the slide. Using needle/brush/forceps, transfer the stained specimen to be mounted in the drop of water. Now, add a drop of glycerine (15% glycerine). Place the coverslip on the glass slide so that one




Fig. 4 How to find out the middle of the slide




Fig. 5a and b How to put coverslip over the sample

of its side touches the liquid on the slide. Hold the other edge with the help of a needle held obliquely (see Fig. 5a). Now, gently lower the needle till the coverslip completely touches the slide. Cover the specimen. Remove the extra fluid from the sides of the coverslip with help of a dropper or a piece of filter paper. Now, the temporary wet mount is ready for observation under microscope.

- c. *Important points for spotting*: Note the spot number and always write it down. The diagram of the spot should be correctly drawn and labelled and write important features of the specimen.

MICROSCOPE

Microscope is a sensitive instrument used to observe things in a laboratory, such as living/dead organism cells and tissues, which cannot be seen by the naked eye.

Magnifying Glass (hand lens)

The simplest type of microscope is the simple magnifying glass (hand lens). It consists of a double convex lens, fitted in a plastic/metal frame, with a handle. It is based on the principle that when an object is placed between a convex lens and its principal focus, an enlarged virtual image is formed. Ordinarily, it magnifies specimens up to 4–5 times depending upon magnification of the lens (3X, 4X, 5X, etc). *It is used to observe specimens, such as an ant, housefly, etc. which do not require higher magnification.*




Fig. 6 Parts of a hand lens

Dissecting Microscope

A dissecting microscope has the following parts:

1. A metallic heavy *base* to support other parts.
2. The *stand* is a short, hollow, cylindrical metallic rod, fixed at one end with the base. It is used for holding the dissecting microscope.
3. *Vertical limb* is a short hollow cylindrical rod which fits into the hollow tube of the stand. An adjustment screw allows it to move up and down to focus an object.
4. The folded arm is a horizontal flat arm attached to the end of the vertical limb, and is used to focus an object by moving it sideways, or up and down. It has a lens fitted at the other end, which may be of magnification 5X, 10X or 20X.
5. *Stage* is a rectangular glass plate attached to the stand. It is used to place the specimen/temporary/permanent slide. It is fitted with two clips to hold the object in place.
6. A *concave mirror* is attached at the lower end of the stand, which helps in reflecting light towards the stage in order to illuminate the object.




Fig. 7 Parts of a dissecting microscope

The magnification of dissecting microscope usually ranges from 5 to 50 times depending upon the power of

the lens, which is usually 5X, 10X or 20X. The dissecting microscope is used to observe small specimens such as mosquito, parts of a small flower, and small organs of specimens. You will also be able to observe different cells in a section of dicot stem.

Compound Microscope

A compound microscope consists of two lenses, called the objective lens and the eyepiece lens, which are placed at proper distances from each other. In a compound microscope, the object to be examined is placed in front of the objective lens, and forms a magnified image. This image is further magnified by the eyepiece lens and made visible to the eye.

Magnification power of a compound microscope depends upon power of objective lens and eyepiece lens. Power of a lens is expressed by the symbol X. It denotes number of times a lens magnifies an object. Generally, a compound microscope consists of eyepiece lens of 10X or 15X magnification. The objective lens are of 10X (*low power*) and 45X (*high power*) magnification.

For *low power magnification* of compound microscope,

$$M = \text{Power of eyepiece lens} \times \text{Power of objective lens} = 10X \times 10X = 100X$$

Thus, under a low power microscope, which has eyepiece lens 10X and objective lens 10X, a specimen can be seen magnified 100 times its original size.

For *high power magnification* of compound microscope,

$$M = \text{Power of eyepiece lens} \times \text{Power of objective lens} = 10X \times 45X = 450X$$

Hence, under high power microscope, a specimen can be seen as magnified as 450 times its size.

A compound microscope consists of the following parts:

1. *Base or Foot* provides stability to the microscope and supports other parts.
2. *Limb or Pillar* is fixed to the foot with a hinge.
3. *Arm* is a curved metal piece which supports the body tube and coarse adjustment.
4. *Inclination joint*, where the arm is attached to the pillar, permits tilting of the microscope to adjust to eye level.
5. *Stage* is a rectangular flat metallic plate, fixed to the lower end of the arm. An opening in its centre admits light to illuminate the object. Two clips are provided over it to hold the slide.
6. *Diaphragm* is present below the stage of the microscope for regulating the amount of light. The diaphragm may be *disc* or *iris type*.
7. *Body tube* is a hollow tube, vertically mounted over the stage, and attached to the upper part of the arm. It can be moved up and down with the help of the screw. The body tube consists of an eyepiece, nose piece and objective lenses.
8. A *coarse adjustment screw* is provided, which can move the body tube up and down for coarse focussing of the object.
9. *Fine adjustment screw* is provided, which can move the body tube very slowly to achieve a fine and sharp focus.
10. A *mirror* is attached below the stage, at the lower end of the arm, to reflect light through the aperture/hole of the stage to the specimen slide.

Setting a Compound Microscope

1. Place the microscope gently on a working/lab table, so that the base is at least 10 cm inside the edge of the table. Keep the arms facing towards yourself.

2. Raise the body tube about 4–5 cm above the stage, by using the coarse adjustment screw.
3. Hold the nose piece and rotate it gently till the low power objective (10X) comes in line with body tube. You will hear a click when this happens. Do not rotate further after the click. Maintain this position even when changing from one objective to another.
4. Slowly, open the diaphragm of the condenser. With one eye, look through the eyepiece lens. Hold the edge of the mirror below the stage, and tilt it towards a light source. When the mirror is in proper position, the circular field in the microscope will be well illuminated.

Focussing the Compound Microscope

1. Turn the low power objective lens of the microscope into position.
2. Place the specimen on the stage, so that the specimen is directly below the objective lens.
3. By using the coarse adjustment screw, lower the body tube till the distance between the objective lens and the cover glass is about 1 mm.
4. Look through the eyepiece lens with one eye, keeping the other eye open. Using the coarse adjustment screw, raise the body tube slowly until the structure of the specimen comes into a clear view. Now, use the fine adjustment and focus to get a sharp image.

Precautions to be taken while using a microscope

1. Always keep the microscope in upright position by holding it with both the hands. Do not swing the microscope with one hand while carrying it, as the lens if loosely fitted, may fall.
2. Always clean the lens, mirror and stage of the microscope properly before use.
3. Do not touch the lens or mirror, as fingerprints may spoil the image.
4. Do not move the lens too low so that it touches the specimen or water.




Fig. 8 Parts of a compound microscope

IN THE CHEMISTRY LABORATORY

How to clean the glass apparatus

Dirty apparatus can give wrong results in an experiment. All glass apparatus should be cleaned and dried well. Usually, a dilute solution of a common detergent is used to wash the apparatus. This should be done under running tap water. You may use bottle brushes if required. Put the apparatus upside-down to dry. Do not dry using a cloth, cotton wool, etc.

If the apparatus is very dirty, the lab assistant can prepare chromic acid solution, by dissolving 5 g of potassium dichromate in 5 mL of water, and adding to it 50 mL of sulphuric acid.

NOTE: This type of wash is done by lab assistant and not by students

Precautions to be taken in a chemistry lab

1. Keep all windows and doors open.
2. The mouth of the test tube should not be pointed towards any pupil including yourself while heating or adding a reagent.



Fig. 9 Correct method of heating a boiling tube



Fig. 10 Wafting technique to know the odour of gas

3. For smelling the vapours, fan the vapours gently with your hands towards the nose. Avoid direct smelling of chemicals or vapours.
4. Acid is diluted by adding water. Never add water to acid.
5. Do not handle chemicals with hands and do not taste any chemical.
6. Do not heat alcohol or any inflammable material directly on flame.
7. Never forcefully insert a thermometer or a glass tube through a cork.




Fig. 11 Correct way of adding acid to water for dilution




Fig. 12 Correct way of inserting a glass tube in a cork





Fig. 13 Some of the common laboratory equipment and glassware

CBSE Living Science Lab Manual Class X



Publisher : **Ratna Sagar**

ISBN : **9789350361528**

Author : **Hiti Mahindroo**

Type the URL : <http://www.kopykitab.com/product/10524>



Get this eBook